

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

12

EUROPEAN PATENT APPLICATION

Application number: 89120001.6

Int. Cl.⁵: C07D 215/227 , A61K 31/47

Date of filing: 27.10.89

Claim for the following Contracting State: ES

Priority: 31.10.88 JP 276953/88

Date of publication of application:
09.05.90 Bulletin 90/19

Designated Contracting States:
CH DE ES FR GB IT LI NL SE

Applicant: OTSUKA PHARMACEUTICAL CO.,
LTD.
9, Kandatsukasa-cho 2-chome
Chiyoda-ku Tokyo(JP)


Inventor: Oshiro, Yasuo

61-11, Yoshinohoncho-6-chome
Tokushima-shi(JP)
Inventor: Sato, Seiji
57-1-A402, Yagami Sasagino, Matsushigecho
Itano-gun Tokushima-ken(JP)
Inventor: Kurahashi, Nobuyuki
463-10, Kagasuno Kawauchicho
Tokushima-shi(JP)


Representative: Kolb, Helga, Dr. Dipl.-Chem. et
al
Hoffmann, Eitle & Partner Patentanwälte
Arabellastrasse 4
D-8000 München 81(DE)

Carbostyryl derivatives.


A novel carbostyryl derivative and salt thereof represented by the formula (1).



(1)




(wherein R is a group of the formula




((wherein R¹ is a C₁-C₃ alkoxy group)), a group of the formula

EP 0 367 141 A2



((wherein R² and R³ are each, at the same time, a chlorine atom, a bromine atom; and R⁴ is a hydrogen atom or a chlorine atom)), 2-methyl-3-nitrophenyl group, 3,5-dichlorophenyl group, or a group of the formula



((wherein R⁵ is a chlorine atom or a bromine atom; and R⁶ is a methyl group)); the carbon-carbon bond between 3- and 4-position in the carbostyryl skeleton is a single or double bond).

The novel carbostyryl derivative and salt thereof represented by the formula (1) is useful agent for treating schizophrenia.

Carbostyryl Derivatives

FIELD OF THE INVENTION

The present invention relates to novel carbostyryl derivatives. More particularly, the invention relates to novel carbostyryl derivatives and salts thereof, processes for preparing said carbostyryl derivatives and salts thereof, as well as pharmaceutical compositions for treating schizophrenia containing, as the active ingredient, said carbostyryl derivative or salt thereof.

BACKGROUND OF THE INVENTION

Schizophrenia is the most common type of psychosis caused by an excessive neurotransmission activity of the dopaminergic nervous system in the central nervous system. [Cf. "Hypothesis of Excessive Dopamine" by Michio Tohr: TAISHA (Metabolism), Vol. 22, pp. 49. (1985); and Pharmacia Review, No. 10, "KOKORO-TO-KUSURI (Mind and Drugs)" edited by Pharmaceutical Society of Japan].

Heretofore, a number of drugs, having the activity for blocking the neurotransmission of dopaminergic receptor in the central nervous system, have been developed, the example for said drugs are phenothiazine-type compounds such as Chlorpromazine; butyrophenone-type compounds such as Haloperidol; and benzamide-type compounds such as Sulpiride. These known drugs are now used widely for the purpose of improving so-called positive symptoms in the acute period of schizophrenia such as hallucinations, delusions and excitations and the like.

However, many of these drugs are considered as not effective for improving so-called the negative symptoms which are observed in the chronic period of schizophrenia such as apathy, emotional depression, hypopsychosis and the like. In addition to the above, these drugs give important side-effects such as akathisia, dystonia, Parkinsonism dyskinesia and late dyskinesia and the like, which are caused by blocking the neurotransmission of dopaminergic receptor in the striate body. Furthermore, other side-effects such as hyperprolactinemia and the like given by these drugs are become other problems. [Cf. G. M. Simpson, E. H. Pi, and J. J. Sramek, Jr.: Drugs, Vol. 21, pp. 138 (1981).]

Under these circumstances, development of drugs for treating schizophrenia having safety and clinically effectiveness have been eagerly expected.

The present inventors have made an extensive study for the purpose of developing drugs for treating schizophrenia, which would be not only effective for improving the negative symptoms, but also effective for improving the positive symptoms of schizophrenia, furthermore such drugs would have less side-effects as compared with those shown by drugs known in prior art. As the result, the present inventors have successfully found carbostyryl derivatives having strong activity for blocking neurotransmission of dopaminergic receptor. As to the side-effects given by known drugs for treating schizophrenia are for example, in the case of phenothiazine-type drugs, the orthostatic hypotension and hypersedation on the basis of strong α -blocking activity; and in the case of drugs having strong activity for blocking neurotransmission of dopaminergic receptor, the side-effects are so-called extrapyramidal tract syndromes such as catalepsy, akathisia, dystonia and the like caused by the blocking neurotransmission of dopaminergic receptor in the striate body.

Among carbostyryl derivatives known in prior art, those disclosed in U. S. Patent No. 4,734,416; Canadian Patent No. 1,117,110; British Patent No. 2,017,701; German Patent Nos. 2,911,108, 1,912,105 and 2,953,723; Japanese Patent Kokai (Laid-open) Nos. 54-130,587 (1979), 55-127,371, (1980) and 62-149,664 (1987) are having chemical structural formulas of upper conception of carbostyryl derivatives of the present invention.

Furthermore, carbostyryl derivatives disclosed in U. S. Patent No. 4,234,585 and European Patent No. 226,441 have chemical structural formula similar to that of carbostyryl derivatives of the present invention, but the pharmacological activities thereof are different from those of possessed by the carbostyryl derivatives of the present invention.

In addition to the above, the carbostyryl derivatives disclosed in U. S. Patent No. 4,234,584 have chemical structural formula similar to that of carbostyryl derivatives of the present invention and also have pharmacological activities similar to those of shown by carbostyryl derivatives of the present invention.

Carbostyryl derivatives disclosed in Australian Patent No. 50252/85, Japanese Patent Kokai (Laid-open) Nos. 58-43952 (1983), 56-49359 (1981), 56-49360 (1981) and 56-49361 (1981) have substituents different

from those of the carbostyryl derivatives of the present invention.

SUMMARY OF THE INVENTION


It is an object of the present invention to provide novel carbostyryl derivatives and salts thereof.

A further object of the present invention is to provide processes for preparing said carbostyryl derivatives and salts thereof.


A still further object of the present invention is to provide a pharmaceutical composition for treating schizophrenia.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS


Carbostyryl derivatives of the present invention and salts thereof are represented by the general formula (1) as follows:




(wherein R is a group of the formula



((wherein R¹ is a C₁-C₃ alkoxy group)), a group of the formula



((wherein R² and R³ are each, at the same time, a chlorine atom, a bromine atom; and R⁴ is a hydrogen atom or a chlorine atom)), 2-methyl-3-nitrophenyl group, 3,5-dichlorophenyl group, or a group of the formula




((wherein R^5 is a chlorine atom or a bromine atom; and R^6 is a methyl group)); the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is a single or double bond), and salts thereof.

Carbostyryl derivatives and salts thereof represented by the general formula (1) possess strong activity for blocking the neurotransmission effect of dopaminergic receptor, with a weak α -blocking activity which have been found during the step of research and development of a number of carbostyryl derivatives, thus when the strength of α -blocking activity of a carbostyryl derivative is defined as the dose (ED_{50} , mg/kg, per os) which is required to inhibits 50% of death of mice being administered with epinephrine, and also the strength of activity for blocking the neurotransmission effect of dopaminergic receptor which is the main activity of carbostyryl derivative, is defined as the dose (ED_{50} mg/kg, per os) which is required to inhibits 50% of stereotypy of mice induced by administration with apomorphine, the agonist of dopamine. The present invention was successfully completed by the above-mentioned findings of said activity.

Carbostyryl derivatives represented by the general formula (1) can be prepared by various methods, the examples for said methods are as follows:


Reaction Formula-1



(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above; and X^1 is a halogen atom or a group which can carry out a substitution reaction similar to a halogen atom, the examples of such group is a mesityloxy group and tosyloxy group and the like).

The reaction of a compound of the general formula (3) with a compound of the general formula (4) can be carried out in the absence or presence of a common inert solvent, under temperature condition of room temperature to 200°C , preferably at 60 to 120°C , and the reaction is completed in about several hours to 24 hours. As to the inert solvent used in this reaction, any solvents for example, ethers such as dioxane, tetrahydrofuran, ethylene glycol dimethyl ether and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; lower alcohols such as methanol, ethanol, isopropanol and the like; polar solvents such as dimethylformamide (DMF), dimethyl sulfoxide (DMSO), acetonitrile and the like can be used. The reaction can be advantageously carried out by using a basic compound as the dehydrohalogenating agent. As to said basic compound, an inorganic basic compound such as calcium carbonate, sodium carbonate, sodium hydroxide, sodium hydrogen carbonate, sodium amide, sodium hydride and the like; and an organic basic compound such as triethylamine, tripropylamine, pyridine, quinoline and the like can be used. Furthermore, the above-mentioned reaction can be carried out, if necessary, by adding an alkali metal iodide such as potassium iodide, sodium iodide or the like as the reaction accelerator. In the above-mentioned reaction, the ratio of used amount of a compound of the general formula (3) to a compound of the general formula (4) may be an equimolar quantity or more, preferably an equimolar quantity to 5 times the molar quantity, more preferably, an equimolar quantity to 1.2 times the molar quantity of the latter to the former.


Reaction Formula-2



15 (wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above; and X² is a halogen atom).

In the Reaction Formula-2, the reaction of a compound represented by the general formula (5) with a compound as the dehydrohalogenating agent, in a suitable solvent at room temperature to 200°C, preferably at 50 to 150°C for within several hours to 15 hours. As to the suitable solvent used in the above reaction, lower alcohols such as methanol, ethanol, isopropanol and the like; ketones such as acetone, methyl ethyl ketone and the like; ethers such as dioxane, diethylene glycol dimethyl ether and the like; aromatic hydrocarbons such as toluene, xylene and the like; DMF, DMSO, hexamethylphosphoryl triamide and the like can be exemplified. As to the basic compound to be used as the dehydrohalogenating agent, an inorganic basic substance such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydride, metallic potassium, sodium amide and the like; an alkali metal alcoholate such as sodium methoxide, sodium ethoxide, potassium ethoxide and the like; as well as an organic basic compound for example, tertiary amines such as pyridine, quinoline, triethylamine, tripropylamine and the like can be exemplified. Furthermore, the above-mentioned reaction can be carried out by using an alkali metal iodide such as potassium iodide, sodium iodide and the like as the reaction accelerator. The ratio of used amount of a compound of the formula (5) to compound of the formula (6) is not specifically restricted, and an equimolar quantity or more of the latter, generally an equimolar to 5 times the molar quantity, preferably an equimolar to 1.2 times of the molar quantity of the latter may be used to one molar quantity of the former.

Reaction Formula-3




50 (wherein R, X² and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above).

The reaction of a compound of the general formula (7) with a compound of the general formula (8) is carried out in a suitable solvent, and in the absence or presence of a basic compound. As to the solvent used in this reaction, aromatic hydrocarbons such as benzene, toluene, xylene and the like; lower alcohols such as methanol, ethanol, propanol, butanol and the like; pyridine, acetone, DMF, DMSO, hexamethylphosphoryl triamide and the like can be exemplified. As to the basic compound used in this reaction, inorganic basic compounds such as sodium carbonate, potassium carbonate, sodium hydrogen carbonate,

potassium hydrogen carbonate, sodium hydroxide, potassium hydroxide, sodium hydride, potassium hydride and the like; organic basic compounds such as triethylamine and the like can be exemplified. A compound of the general formula (8) may be used at least an equimolar quantity, preferably an equimolar to 3 times the molar quantity thereof to one molar quantity of a compound of the general formula (7). The reaction is carried out, generally at room temperature to 180°C, preferably at 80 to 150°C, and is completed in about 3 to 30 hours.

Reaction Formula-4




(wherein R, X' and the carbon-carbon bond between 3- and 4- positions in the carbostyryl skeleton are the same as defined above).

The reaction of a compound of the formula (9) with a compound of the formula (10) can be carried out in a suitable solvent and in the absence or presence of a basic compound. As to the solvent used in this reaction, water; a lower alcohols such as methanol, ethanol, isopropanol, butanol and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; acetic acid, ethyl acetate, DMF, DMSO, hexamethylphosphoryl triamide and the like can be exemplified. As to the basic compound used in this reaction, an inorganic basic compound such as sodium carbonate, potassium carbonate, sodium hydrogen carbonate, sodium hydroxide, potassium hydroxide and the like; an alkali metal alcoholate such as sodium methylate, sodium ethylate and the like; an organic basic compound such as 1,5-diazabicyclo[4.3.0]nonene-5 (DBN), 1,8-diazabicyclo[5.4.0]undecene-7 (DBU), 1,4-diazabicyclo[2.2.2]octane (DABCO) and the like can be exemplified.

A compound of the general formula (10) may be used generally, at least an equimolar quantity, preferably an equimolar to 5 times the molar quantity to one molar quantity of compound of the general formula (9). The reaction is generally carried out at 40 to 120°C, preferably at about 70 to 100°C, and is completed in about 1 to 15 hours.

Reaction Formula-5




(wherein R, X' and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above).

The reaction of a compound of the general formula (11) with a compound of the general formula (12) is carried out under conditions similar to those employed in the reaction of a compound of the general formula (9) with a compound of the general formula (10).

In the above-mentioned Reaction Formula-1, a compound of the general formula (4) used as one of the starting materials is prepared by a method as shown in Reaction Formula-6 as follows.

Reaction Formula-6



(wherein R and X¹ are the same as defined above).

The reaction of a compound of the formula (10) with a compound of the formula (13) is carried out by methods similar to those employed in the reaction of a compound of the formula (9) with a compound of the formula (10).

Carbostyryl derivative represented by the formula (1) of the present invention can easily be converted into its acid-addition salt by reacting it with a pharmaceutically acceptable acid. The examples of such acid includes inorganic acids such as hydrochloric acid, sulfuric acid, phosphoric acid, hydrobromic acid and the like; organic acids such as oxalic acid, maleic acid, fumaric acid, malic acid, tartaric acid, citric acid, benzoic acid and the like. Among carbostyryl derivatives represented by the formula (1) of the present invention, those having acidic group can easily be converted into their salts by reacting with basic compounds. The examples of such basic compounds includes sodium hydroxide, potassium hydroxide, calcium hydroxide, sodium carbonate, potassium hydrogen carbonate and the like. The desired compounds prepared by the procedures in the above-mentioned various reaction formulas can easily be isolated and purified by usual separation means such as solvent extraction, dilution, recrystallization, column chromatography, preparative thin layer chromatography and the like.

Carbostyryl derivatives represented by the general formula (1) can be used in the form of usual pharmaceutical compositions which are prepared by using diluents or excipients such as fillers, bulking agents, binders, wetting agents, disintegrating agents, surface active agents, lubricants and the like. As to the pharmaceutical compositions, various types of administration unit forms can be selected depending on the therapeutical purposes, and the examples of pharmaceutical compositions are tablets, pills, powders, liquids, suspensions, emulsions, granules, capsules, suppositories, injection preparations (solutions and suspensions) and the like. For the purpose of shaping the pharmaceutical composition in the form of tablets, any excipients which are known and used widely in this field can also be used, for example carriers such as lactose, white sugar, sodium chloride, glucose, urea, starch, calcium carbonate, kaolin, crystalline cellulose, silicic acid and the like; binders such as water, ethanol, propanol, simple sirup, glucose solutions, starch solutions, gelatin solutions, carboxymethyl cellulose, shalac, methyl cellulose, potassium phosphate, polyvinylpyrrolidone and the like; disintegrating agents such as dried starch, sodium alginate, agar powder, laminaria powder, sodium hydrogen carbonate, calcium carbonate, fatty acid esters of polyoxyethylene sorbitan, sodium laurylsulfate, monoglyceride of stearic acid, starch, lactose and the like; disintegration inhibitors such as white sugar, stearin, coconut butter, hydrogenated oils; absorption accelerators such as quaternary ammonium base, sodium laurylsulfate and the like; wetting agents such as glycerin, starch and the like; adsorbing agents such as starch, lactose, kaolin, bentonite, colloidal silicic acid and the like; and lubricants such as purified talc, stearates, boric acid powder, polyethylene glycol and the like. If tablets are desired, they can be further coated with the usual coating materials to make the tablets as sugar coated tablets, gelatin film coated tablets, tablets coated with enteric coatings, tablets coated with films, double layered tablets and multi-layered tablets.

For the purpose of shaping the pharmaceutical composition in the form of pills, any excipients which are known and widely used in this field can also be used, for example, carriers such as lactose, starch, coconut butter, hardened vegetable oils, kaolin, talc and the like; binders such as gum arabi powder, tragacanth gum powder, gelatin, ethanol and the like; disintegrating agents such as agar, laminaria and the like.

For the purpose of shaping the pharmaceutical composition in the form of suppositories, any excipients

which are known and widely used in this field can also be used, for example polyethylene glycols, coconut butter, higher alcohols, esters of higher alcohols, gelatin, semi-synthesized glycerides and the like.

For the purpose of shaping the pharmaceutical composition in the form of injection preparations, solutions and suspensions are sterilized and are preferably made isotonic to blood. In making injection preparations, any carriers which are usually used in this field can also be used, for example, water, ethyl alcohol, propylene glycol, ethoxylated isostearyl alcohol, polyoxylated isostearyl alcohol, fatty acid esters of polyoxyethylene sorbitan. In these instances, adequate amounts of sodium chloride, glucose or glycerin can be added to the desired injection preparations to make them isotonic. Furthermore, usual dissolving agents, buffer agents, analgesic agents may be added. Yet further, if necessary, coloring agents, preservatives, perfumes, seasoning agents, sweetening agents and other medicines may also be added to the desired preparations during the treatment of schizophrenia.

The amount of carbostyryl derivative of the general formula (1) or salt thereof to be contained in a pharmaceutical composition for treating schizophrenia according to the present invention is not specifically restricted and can suitably be selected from a wide range, usually it is contained 1 to 70%, preferably 1 to 30% by weight of the whole composition.

Administration methods of a pharmaceutical composition for treating schizophrenia of the present invention are not specifically restricted, and can be administered in various forms of preparations depending on the age of the patient, distinction of sex, other conditions, as well as conditions of the symptoms. For example, tablets, pills, solutions, suspensions, emulsions, granules and capsules are orally administered; and injection preparations are administered singly or mixed with injection transfusions such as glucose solutions and amino acid solutions intravenously; and if necessary, the injection preparations are administered singly intramuscularly, intracutaneously, subcutaneously or intraperitoneally. Suppositories are administered into the rectum.

The dosage of a pharmaceutical composition for treating schizophrenia according to the present invention are suitably selected according to the method of use, the age of the patient, distinction of sex, other conditions, as well as conditions of the symptoms, usually about 0.1 to 10 mg/kg of the body weight/day of carbostyryl derivative of the general formula (1) as the active ingredient may be administered. Usually, 1 to 200 mg of the active ingredient may be contained in an administration unit form.

In the above-mentioned formula (1), the C₁-C₃ alkoxy group is a straight-chain or branched-chain alkoxy group having 1 to 3 carbon atoms, such as a methoxy group, an ethoxy group, a propoxy group, an isopropoxy group and the like, and among these, methoxy group and ethoxy group are preferable, and ethoxy group is the most preferable. Furthermore, the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is preferably a single bond.

The present invention will be explained in detail by showing Reference Examples, Examples, Pharmacological test results and Examples of Pharmaceutical Compositions, however, the present invention are not restricted only thereto.

Reference Example 1

To a mixture of 6.08 g of 2-chloro-3-methylaniline, 9 g of di(2-bromoethyl)amine hydrobromide and 4 ml of water was added a solution of 0.8 g of potassium hydroxide and 2.5 ml of water 3 times of 1 hour interval at 100°C, then the reaction mixture was stirred at the same temperature for 9 hours. To the resultant reaction mixture was added potassium hydroxide to make the mixture alkaline, and the mixture was extracted with diethyl ether, washed with water, dried with anhydrous magnesium sulfate. The solvent was removed by evaporation and the residue thus obtained was purified by means of a silica gel column chromatography (eluent: 5%-methanol/chloroform), and obtained 3.41 g of 4-(2-chloro-3-methylphenyl)-piperazine.





Light purple oily substance

¹H-NMR (CDCl₃) δ: 2.38 (3H, s), 3.04 (8H, m), 6.93 (2H, m), 7.12 (1H, dd, J = 7.7Hz, 7.7Hz)

Reference Examples 2-5

By procedures similar to those employed in the above mentioned Reference Example 1, by using suitable starting materials, there were prepared compounds of Reference Examples 2-5 as shown in the following Table 1.

Table 1

Reference Example No.	R	$^1\text{H-NMR}$ (CDCl_3) δ :
2		2.45 (3H, s), 2.90 (4H, m), 3.05 (4H, m), 7.23 (1H, dd, J=8.0Hz, 2.0Hz), 7.28 (1H, dd, J=7.4Hz, 8.0Hz), 7.52 (1H, dd, J=7.4Hz, 2.0Hz)
3		2.42 (3H, s), 3.03 (8H, m), 6.90 (1H, d, J=7.9Hz), 6.95 (1H, d, J=7.5Hz), 7.17 (1H, dd, J=7.5Hz, 7.9Hz)
4		3.05 (8H, m), 6.91 (1H, d, J=2.3Hz), 7.17 (1H, d, J=2.3Hz)
5		3.02 (8H, m), 6.98 (1H, dd, J=8.0Hz, 1.5Hz), 7.14 (1H, t, J=8.0Hz), 7.35 (1H, dd, J=8.0Hz, 1.5Hz)

Reference Example 6

To a solution of 4.06 g of potassium carbonate with 400 ml of water was added 40 g of 7-hydroxy-3,4-dihydrocarbostyryl and 158 g of 1,4-dibromobutane, then the mixture was refluxed for 3 hours. The reaction mixture thus obtained was extracted with dichloromethane, dried with anhydrous magnesium sulfate, then the solvent was removed by evaporation. The residue thus obtained was purified by means of a silica gel column chromatography (eluent: dichloromethane), and recrystallized from n-hexane-ethanol to yield 50 g of 7-(4-bromobutoxy)-3,4-dihydrocarbostyryl.

Coloreless needle crystals
Melting point: 110.5 - 111.0 °C.

5 Example 1

A suspension of 47 g of 7-(4-bromobutoxy)-3,4-dihydrocarbostyryl, 35 g of sodium iodide with 600 ml of acetonitrile was refluxed for 30 minutes. To this suspension was added 40 g of 1-(2,3-dichlorophenyl)-piperazine and 33 ml of triethylamine and the whole mixture was further refluxed for 3 hours. After the
10 solvent was removed by evaporation, the residue thus obtained was dissolved in chloroform, washed with water then dried with anhydrous magnesium sulfate. The solvent was removed by evaporation, and the residue thus obtained was recrystallized from ethanol twice, to yield 57.1 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

Colorless flake crystals

15 Melting point: 139.0 - 139.5 °C.

One gram of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl was dissolved in 20 ml of ethanol by heating, then under stirring condition, an ethanol solution saturated with hydrogen chloride was added thereto, the crystals precipitated were collected by filtration and recrystallized from
20 thanol to yield 0.75 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl hydrochloride.

White powdery substance

Melting point: 214-222 °C. (decomposed).

One gram of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl was dissolved in 10 ml of ethanol, then to this solution was added 4 ml of sulfuric acid-ethanol (1ml of concentrated sulfuric
25 acid/10 ml of ethanol), then the solvent was removed by evaporation. To the residue thus obtained was added 10 ml of ethanol and 30 ml of water, the mixture was heated to make it as a solution, recrystallized, and the crystals were collected by filtration, further recrystallized from ethanol-water to yield 1.02 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl* sulfate.

White powdery substance

30 Melting point: 220-225 °C.

By using 1.0 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl and 290 mg of fumaric acid, and treated by procedures similar to those employed in the case of preparation of the sulfate as mentioned above, and recrystallized from ethanol to yield 0.97 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl* fumarate.

35 White powdery substance

Melting point: 196-198 °C.

By using 1.0 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl and 290 mg of maleic acid, and treated by procedures similar to those employed in the case of preparation of the sulfate as mentioned above, and recrystallized from ethanol to yield 0.98 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl* maleate.
40

White powdery substance

Melting point: 172-180 °C.

45 Examples 2-14

By using suitable starting materials, and by procedures similar to those employed in Example 1, there were prepared compounds of Examples 2-14 as shown in Table 2 as follows. In Table 2, compounds of Examples 11-14 are in the form of hydrochlorides.


50

55

Table 2

5

10



15

Example
No.

R


Carbon-carbon
bond between
3- and 4-
positions in
carbostyryl
skeleton

Crystal form
(Recrystallization
solvent)

Melting
point
(°C)

20

2




Single bond

Yellow needle
crystals
(Methanol)

165-166

25

3




Single bond

Colorless flake
crystals
(Ethanol)

133-134

30

4



Single bond

Colorless needle
crystals
(Ethanol)

125-126

35

(To be continued)

40

45

50





55

Table 2 (Continued)

Example No.	R	Carbon-carbon bond between 3- and 4-positions in carbostyryl skeleton	Crystal form (Recrystallization solvent)	Melting point (°C)
5		Single bond	White powdery substance (Ethanol)	134-135
6		Single bond	Colorless granular crystals (Ethanol)	133-134
7		Single bond	White powdery substance (Methanol)	174-176
8		Single bond	White powdery substance (Methnaol)	125-126
9		Single bond	Pale brown flake crystals (Methnaol)	150-151
10		Double bond	White powdery substance (Ethanol)	144-146

(To be continued)

Table 2 (Continued)

Example No.	R	Carbon-carbon bond between 3- and 4-positions in carbostyryl skeleton	Crystal form (Recrystallization solvent)	Melting point (°C)
11		Double bond	White powdery substance (Ethanol)	151 (decomp.)
12		Single bond	Colorless fine needle crystals (Ethanol)	214-218
13		Single bond	Pale brown powdery substance (Ethanol-diethyl ether)	207-207.5
14		Single bond	Pale brown powdery substance (Ethanol-diethyl ether)	203-203.5

40 Pharmacological Tests

a) Anti-apomorphine activity in mouse:

45 Pharmacological test was conducted by using six mice in one test group. One hour after the oral administration of a test compound to a test mouse, apomorphine (1.25 mg/kg) was subcutaneously administered, and the stereotypy movements manifested were scored according to the method by Puech (Neuropharmacology, Vol. 20, pp. 1279, 1981). The anti-apomorphine activity performed by each of the test compounds were evaluated by the scored data as the indication thereof.

50 50% Effective dose (ED₅₀, mg/kg) of anti-apomorphine activity performed by a test compound is determined in that when the score obtained from the test group is lower than 50% of mean value of the score obtained from the control group, then it is defined as "positive" in anti-apomorphine activity.

55 b) Anti-epinephrine lethal activity in mouse:

By procedures similar to those described in Janssen, P., et al.: Arzneimittel Forschung, Vol. 13, pp. 205, (1963), the test was conducted by using six mice in one test group. One hour after the oral

administration of a test compound to a test mouse, lethal dose (1.5 mg/kg) of epinephrine was intravenously administered, and 4 hours after of the intravenous administration, each of mice in the test group was observed whether it is

50% Effective dose (ED₅₀ mg/kg) of anti-epinephrine lethal activity performed by a test compound is determined from the amount thereof orally administered, and in the case that the mouse is alived is determined as "positive" in anti-epinephrine lethal activity.

The test results are shown in Table 3 as follows.

Test compound No.	
1	Compound of Example 1 (Free form)
2	Compound of Example 2
3	Compound of Example 3
4	Compound of Example 4
5	Compound of Example 5
6	Compound of Example 12
7	Compound of Example 7
8	Compound of Example 8
9	Compound of Example 9
10	Compound of Example 10
11	Compound of Example 11
12	Compound of Example 13
13	Compound of Example 14

Table 3

	Anti-apomorphine activity	Anti-epinephrine activity	
Test compound No.	(ED ₅₀ mg/kg) (A)	(ED ₅₀ mg/kg) (B)	(B)/(A)
1	0.18	>128	>711
2	0.3	>128	>426.7
3	0.4	> 64	>160
4	0.4	> 64	>160
5	0.5	>128	>256
6	0.1	3.7	37
7	0.4	>128	>320
8	0.2	2.5	12.5
9	0.6	>256	>426.7
10	0.36	>128	>355
11	0.12	3.8	31.6
12	0.5	1.58	3.16
13	0.2	0.24	1.2

Example of Preparation of Pharmaceutical Composition -1		
7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl		5 mg
Starch		132 mg
Magnesium stearate		18 mg
Lactose		45 mg
Total	Total	200 mg



By using usual procedures, tablets containing the above formulation per one tablet were prepared.

Example of Preparation of Pharmaceutical Composition-2		
7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl		500 mg
Polyethylene glycol (Molecular weight: 4,000)		0.3 g
Sodium chloride		0.9 g
Polyoxyethylene sorbitan monooleate		0.4 g
Sodium metabisulfite		0.1 g
Methyl p-hydroxybenzoate		0.18 g
Propyl p-hydroxybenzoate		0.02 g
Distilled water for injection		100 ml


The above-mentioned methyl p-hydroxybenzoate, propyl p-hydroxybenzoate, sodium metabisulfite and sodium chloride were dissolved in distilled water for injection at 80° C with stirring. The resulting solution was cooled to 40° C, then 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl, polyethylene glycol and polyoxyethylene sorbitan monooleate were dissolved in the above-mentioned solution in this order, then the predetermined volume of the injection solution was adjusted by adding the distilled water for injection, and was sterilized by filtration by using a suitable filter paper, then 1 ml each of the desired injection solution was filled in an ampul.

Claims


1. A carbostyryl derivative and salt thereof represented by the formula (1),




(wherein R is a group of the formula



((wherein R¹ is a C₁-C₃ alkoxy group)), a group of the formula




((wherein R² and R³ are each, at the same time, a chlorine atom, a bromine atom; and R⁴ is a hydrogen atom or a chlorine atom)), 2-methyl-3-nitrophenyl group, 3,5-dichlorophenyl group, or a group of the formula




((wherein R⁵ is a chlorine atom or a bromine atom; and R⁶ is a methyl group)); the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is a single or double bond), and salt thereof.

2. The carbostyryl derivative and salt thereof according to Claim 1, wherein R is a group of the formula



(wherein R¹ is the same as defined above).


3. The carbostyryl derivative and salt thereof according to Claim 1, wherein R is a group of the formula



(wherein R², R³ and R⁴ are the same as defined above).

4. The carbostyryl derivative and salt thereof according to Claim 1, wherein R is a 2-methyl-3-nitrophenyl group or a 3,5-dichlorophenyl group.

5. The carbostyryl derivative and salt thereof according to Claim 1, wherein R is a group of the formula



(wherein R^5 and R^6 are the same as defined above).

6. The carbostyryl derivative and salt thereof according to Claim 2, wherein R^1 is an ethoxy group.

7. The carbostyryl derivative and salt thereof according to Claim 3, wherein R^2 and R^3 are, at the same time, chlorine atoms, and R^4 is a hydrogen atom.

8. The carbostyryl derivative and salt thereof according to Claim 3, wherein R^2 and R^3 are, at the same time, bromine atoms, and R^4 is a hydrogen atom.

9. The carbostyryl derivative and salt thereof according to Claim 3, wherein R^2 , R^3 and R^4 are, at the same time, chlorine atoms.

10. The carbostyryl derivative and salt thereof according to Claim 5, wherein R^5 is a chlorine atom.

11. The carbostyryl derivative and salt thereof according to Claim 5, wherein R^5 is a bromine atom.


12. 7-{4-[4-(2,3-Dichlorophenyl)-1-piperazinyl]-butoxy}-3,4-dihydrocarbostyryl.

13. 7-{4-[4-(2,3-Dichlorophenyl)-1-piperazinyl]-butoxy}carbostyryl.


14. 7-{4-[4-(2-Ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

15. 7-{4-[4-(2-Ethoxyphenyl)-1-piperazinyl]butoxy}-carbostyryl.


16. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1).




(wherein R is a group of the formula



((wherein R^1 is a C_1 - C_3 alkoxy group)), a group of the formula



((wherein R^2 and R^3 are each, at the same time, a chlorine atom, a bromine atom; and R^4 is a hydrogen atom or a chlorine atom)), 2-methyl-3-nitrophenyl group, 3,5-dichlorophenyl group, or a group of the formula




5

((wherein R^5 is a chlorine atom or a bromine atom; and R^6 is a methyl group)); the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is a single or double bond) by reacting a carbostyryl compound of the formula (3)

10

15




(3)

20

(wherein X^1 is a halogen atom or a group which can carry out a substitution reaction similar to a halogen atom; and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above) with a piperazine compound of the formula (4).

25




(4)

(wherein R is the same as defined above).

30

17. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),

35




(1)

40

(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above) by reacting a carbostyryl compound of the formula (5),

45


50



(5)


(wherein the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above) with a piperazine compound of the formula (6),

55




(wherein R is the same as defined above; and X^2 is a halogen atom).

18. A process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),



(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above)

by reacting a carbostyryl compound of the formula (7),




(wherein the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above) with a compound of the formula (8),

$R - X^2$ (8)

(wherein R and X^2 are the same as defined above).

19. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),




(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above),

by reacting a carbostyryl compound of the formula (9),

5

10



(wherein X¹ and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above),

15 with a compound of the formula (10),


NH₂-R (10)

(wherein R is the same as defined above).

20. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),

20

25




30

(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above),

by reacting a carbostyryl compound of the formula (11),


35

40



45 (wherein the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above), with a compound of the formula (12),

50



55 (wherein X¹ and R are the same as defined above).

21. A pharmaceutical composition containing, as the active ingredient, a carbostyryl derivative or salt thereof represented by the formula (1) as claimed in Claim 1 and pharmaceutically acceptable carriers.

22. The pharmaceutical composition according to Claim 21, wherein the carbostyryl derivative or salt

thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

23. The pharmaceutical composition according to Claim 21, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}carbostyryl.

24. The pharmaceutical composition according to Claim 21, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

25. The pharmaceutical composition according to Claim 21, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}carbostyryl.

26. A pharmaceutical composition for treating schizophrenia containing, as the active ingredient, a carbostyryl derivative or salt thereof represented by the formula (1) as claimed in Claim 1 and pharmaceutically acceptable carriers.

27. The pharmaceutical composition for treating schizophrenia according to Claim 26, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

28. The pharmaceutical composition for treating schizophrenia according to Claim 26, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}carbostyryl.

29. The pharmaceutical composition for treating schizophrenia according to Claim 26, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

30. The pharmaceutical composition for treating schizophrenia according to Claim 26, wherein the carbostyryl derivative or salt thereof is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}carbostyryl.

31. The use of a carbostyryl derivative and salt thereof represented by formula (1) as claimed in claim 1 for the preparation of a drug useful in the treatment of schizophrenia.

32. The use according to claim 31, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.


33. The use according to claim 31, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}carbostyryl.

34. The use according to claim 31, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.


35. The use according to claim 31, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}carbostyryl.

Claims for the following Contracting State: ES


1. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),




(wherein R is a group of the formula




((wherein R¹ is a C₁-C₃ alkoxy group)), a group of the formula



10 ((wherein R^2 and R^3 are each, at the same time, a chlorine atom, a bromine atom; and R^4 is a hydrogen atom or a chlorine atom)), 2-methyl-3-nitrophenyl group, 3,5-dichlorophenyl group, or a group of the formula




20 ((wherein R^5 is a chlorine atom or a bromine atom; and R^6 is a methyl group)); the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is a single or double bond) by reacting a carbostyryl compound of the formula (3)



(3)


35 (wherein X^1 is a halogen atom or a group which can carry out a substitution reaction similar to a halogen atom; and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above) with a piperazine compound of the formula (4),



(4)

(wherein R is the same as defined above).


2. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),



(1)


55 (wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined in claim 1),

by reacting a carbostyryl compound of the formula (5),



(5)


(wherein the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above) with a piperazine compound of the formula (6),



(6)


(wherein R is the same as defined above; and X² is a halogen atom).

3. A process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),



(1)

(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined in claim 1),
by reacting a carbostyryl compound of the formula (7),



(7)


(wherein the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above) with a compound of the formula (8),
R - X² (8)

(wherein R and X² are the same as defined in claims 1 and 2).

4. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),

5

10



(1)


(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined in claim 1),

15

by reacting a carbostyryl compound of the formula (9),

20

25



(9)

(wherein X¹ and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined above),

30

with a compound of the formula (10),

NH₂-R (10)


(wherein R is the same as defined above).

5. Process for preparing a carbostyryl derivative and salt thereof represented by the formula (1),

35

40

45




(1)

(wherein R and the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton are the same as defined in claim 1),

by reacting a carbostyryl compound of the formula (11),


50

55



(11)

(wherein the carbon-carbon bond between 3- and 4-positions in the carbostyryl skeleton is the same as defined above), with a compound of the formula (12),



10 (wherein X¹ and R are the same as defined in claim 1).

6. The process according to any of claims 1 to 5, wherein R is a group of the formula



20 (wherein R¹ is the same as defined in claim 1).

7. The process according to any of claims 1 to 5, wherein R is a group of the formula



(wherein R², R³ and R⁴ are the same as defined in claim 1).

8. The process according to any of claims 1 to 5, wherein R is a 2-methyl-3-nitrophenyl group or a 3,5-dichlorophenyl group.

35 9. The process according to any of claims 1 to 5, wherein R is a group of the formula



45 (wherein R⁵ and R⁶ are the same as defined in claim 1).

10. The process according to any of claims 1 to 5, wherein R¹ is an ethoxy group.

11. The process according to claims 1 to 5, wherein R² and R³ are, at the same time chlorine atoms, and R⁴ is a hydrogen atom.

50 12. The process according to claims 1 to 5, wherein R² and R³ are, at the same time bromine atoms, and R⁴ is a hydrogen.

13. The process according to claims 1 to 5, wherein R², R³ and R⁴ are, at the same time, chlorine atoms.

14. The process according to claims 1 to 5, wherein R⁵ is a chlorine atom.

55 15. The process according to claims 1 to 5, wherein R⁵ is a bromine atom.

16. The process according to claims 13 to 15, wherein the prepared compound is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]-butoxy}-3,4-dihydrocarbostyryl.

17. The process according to claims 13 to 15, wherein the prepared compound is 7-{4-[4-(2,3-

dichlorophenyl)-1-piperazinyl]butoxy}-carbostyryl.

18. The process according to claims 13 to 15, wherein the prepared compound is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

19. The process according to claims 13 to 15, wherein the prepared compound is 7-{4-[4-(2-ethoxyphenyl)-1-piperazinyl]butoxy}-carbostyryl.

20. The use of a carbostyryl derivative and salt thereof as represented by the formula (1) of claim 1 for the preparation of a pharmaceutical.

21. The use of a carbostyryl derivative and salt thereof as represented by the formula (1) of claim 1 for the preparation of a pharmaceutical useful in the treatment of schizophrenia.

22. The use according to claims 20 and 21, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

23. The use according to claims 20 and 21, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-carbostyryl.

24. The use according to claims 20 and 21, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2,3-ethoxyphenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

25. The use according to claims 20 and 21, wherein the carbostyryl derivative and salt thereof is 7-{4-[4-(2,3-ethoxyphenyl)-1-piperazinyl]butoxy}-carbostyryl.

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets




(11) Publication number:

0 367 141 A3


(12)


EUROPEAN PATENT APPLICATION(21) Application number: **89120001.6**(51) Int. Cl.⁵: **C07D 215/227, A61K 31/47**(22) Date of filing: **27.10.89**(30) Priority: **31.10.88 JP 276953/88**(43) Date of publication of application:
09.05.90 Bulletin 90/19(84) Designated Contracting States:
CH DE ES FR GB IT LI NL SE(88) Date of deferred publication of the search report:
12.06.91 Bulletin 91/24(71) Applicant: **OTSUKA PHARMACEUTICAL CO., LTD.**
9, Kandatsukasa-cho 2-chome
Chiyoda-ku Tokyo(JP)(72) Inventor: **Oshiro, Yasuo**
61-11, Yoshinohoncho-6-chome
Tokushima-shi(JP)
Inventor: **Sato, Seiji**
57-1-A402, Yagami Sasagino, Matsushigecho
Itano-gun Tokushima-ken(JP)
Inventor: **Kurahashi, Nobuyuki**
463-10, Kagasuno Kawauchi-cho
Tokushima-shi(JP)(74) Representative: **Kolb, Helga, Dr. Dipl.-Chem. et al**
Hoffmann, Eitle & Partner Patentanwälte
Arabellastrasse 4
W-8000 München 81(DE)(54) **Carbostyryl derivatives.**

(57) A novel carbostyryl derivative and salt thereof represented by the formula (1),




(wherein R is a group of the formula

((wherein R¹ is a C₁-C₃ alkoxy group)), a group of the formula**EP 0 367 141 A3**



((wherein R² and R³ are each, at the same time, a chlorine atom, a bromine atom; and R⁴ is a hydrogen atom or a chlorine atom)), 2-methyl-3-nitrophenyl group, 3,5-dichlorophenyl group, or a group of the formula



((wherein R⁵ is a chlorine atom or a bromine atom; and R⁶ is a methyl group)); the carbon-carbon bond between 3- and 4-position in the carbostyryl skeleton is a single or double bond).

The novel carbostyryl derivative and salt thereof represented by the formula (1) is useful agent for treating schizophrenia.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

Application number

EP 89 12 0001

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	GB-A-2 017 701 (OTSUKA PHARMACEUTICAL CO. LTD) * Claim 1; page 4, line 35; examples 142,205,233,258,294 * & US-A-4 734 416 (Cat. D) --	1,21, 31	C 07 D 215/227 A 61 K 31/47
X	EP-A-0 182 247 (HOECHST AG) * Claim 1; page 8, compounds 1-3,5 * --	1,21, 31	
X	EP-A-0 005 828 (HOECHST AG) * Claims 1,3; examples 35,36 * & US-A-4 234 584 (Cat. D) -- ./.	1,21, 31	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 D 215/00 A 61 K 31/00
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-25, 31-35</p> <p>Claims searched incompletely:</p> <p>Claims not searched: 26-30</p> <p>Reason for the limitation of the search:</p> <p>Method for treatment of the human or animal body by surgery or therapy (See art. 52(4) of the European Patent Convention)</p>			
Place of search THE HAGUE		Date of completion of the search 07-03-1991	Examiner BOSMA
CATEGORY OF CITED DOCUMENTS		<p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>	
<p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p>			

EPO Form 1501.1.03 R2



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application number

EP 89 12 0001

-2-

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	GB-A-2 071 094 (OTSUKA PHARMACEUTICAL CO. LTD) * Claim 1; page 2, lines 46-56 * --	1,21,31	
A	US-A-3 994 900 (J. KRAPCHO et al.) --		
A	EP-A-0 006 506 (BOEHRINGER MANNHEIM GmbH) * Claim 1; example 9 * & US-A-4 234 585 (Cat. D) --	1,21,31	
X,P	CHEMICAL & PHARMACEUTICAL BULLETIN vol. 36, no. 11, November 1988, pages 4377-4388, Tokyo, JP; K. BANNO et al.: "Studies on 2(1H)-Quinolinone derivatives as Neuroleptic agents. I. Synthesis and biological activities of (4-phenyl-1-piperazinyl)-propoxy-2(1H)-quinolinone derivatives" * Table II * --	1,21,31	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
A,D	EP-A-0 226 441 (OTSUKA PHARMACEUTICAL CO. LTD) * Claim 1 * -----	1,21,31	